

Thermodynamics (Classical) for Biological systems

Prof. G. K. Suraishkumar

Department of Biotechnology

Indian Institute of Technology Madras

Module No. # 06

Reaction Equilibria

Lecture No. # 37

Reaction in Liquid or Solid Phases

Welcome!

In the last class we looked at the temperature dependence of the equilibrium constant. We had looked at two equilibrium constants, K_p and K_f . All those were valid for gas phase reactions.

(Refer Slide Time: 00:35)

Equilibrium constant for reactions occurring in liquid/solid solutions

To derive an expression for the equilibrium constant in solutions, let us begin with the same equation as earlier, i.e. Eq. 6.14

$$\sum_i \nu_i \mu_i = 0$$

Substitution of the expression for the relevant chemical potential (Eq. 4.8) in Eq. 6.14 yields

$$\sum_i \nu_i (\mu_i^\# + R T \ln \gamma_i x_i) = 0 \quad \text{Eq. 6.43}$$

NPTEL

... What we are going to start doing in this class is to consider equilibrium constant for reactions that occur in liquid or solid solutions. To derive an expression for the equilibrium constant in solutions, let us begin with the same equation that is the equation 6.14, which is sum over $\nu_i \mu_i$ equals zero. This is the criterion for reaction equilibrium; this is something that we said is very fundamental good to remember and so on. Simple – sum over all ν_i the stoichiometric coefficient times the chemical potential of the various species μ_i equals zero. If, we substitute the expression for the


relevant chemical potential ... this is for liquid or solid solution. Therefore, you know that it is going to be $\mu_i = \mu_i^\# + RT \ln \gamma_i x_i$; that is how we characterize the liquid and solid solutions. If we do that sum over i ν_i and we are going to replace μ_i with $\mu_i^\# + RT \ln \gamma_i x_i$; this equals zero. Let us call this equation 6.43.

(Refer Slide Time: 01:58)

which can be expressed as:

$$\sum_i \nu_i \mu_i^\# + RT \sum_i \nu_i \ln \gamma_i x_i = 0$$

Or,

$$\sum_i \nu_i \mu_i^\# = -RT \ln \prod_i (\gamma_i x_i)^{\nu_i} \quad \text{Eq. 6.44}$$


Which can be expressed as ... I am just going to expand the terms here ... sum over all i $\nu_i \mu_i^\# + RT \sum_i \nu_i \ln \gamma_i x_i$. RT is a constant. Therefore, I have taken it out here. This equals zero. Or you could write this as sum over i $\nu_i \mu_i^\# = -RT \ln \dots$ I have taken the \ln out here; or, before that \ln of $\nu_i \ln$ of $\gamma_i x_i$ which is nothing but \ln of $\gamma_i x_i$ raised to the power ν_i If I take the log out, the sum of the log is nothing but the log of the product of the various things and therefore, log of the product over all i $\gamma_i x_i$ the whole raised to ν_i . We will call it equation 6.44.

(Refer Slide Time: 03:08)


If we define,

$$\prod_i (\gamma_i x_i)^{\nu_i} \equiv K \quad \text{Eq. 6.45}$$

then, we can write Eq. 6.44 as

$$-R T \ln K = \sum_i \nu_i \mu_i^\# \quad \text{Eq. 6.46}$$

Since $\mu_i^\#$ is a function of T and P , K can also be expected to depend both on T and P



Now, if we define the same way that we did for K_p and so on, if we define the product over all i of $\gamma_i x_i$ raised to the power ν_i as a certain K , equation 6.45. This is the equilibrium constant for reactions occurring in solid or liquid phases. Then, we can write our previous equation as $\text{minus } RT \ln K = \text{sum over } i \nu_i \mu_i^\#$. Now, are you able to see the relationship or the similarity between this expression 6.46 and $\text{minus } RT \ln K_p = \text{sum over } i \nu_i \mu_i^\circ$? That was μ_i° , this is $\mu_i^\#$.

There it was a function only of temperature μ_i° , but here since $\mu_i^\#$ is a function of both temperature and pressure, you expect the equilibrium constant, here also to be a function of both temperature and pressure – think, I say that here. Since, $\mu_i^\#$ is a function of temperature and pressure, K can also be expected to depend on both the temperature and pressure.

(Refer Slide Time: 04:27)


Eq. 6.46 can be rearranged as:

$$R \ln K = - \frac{\sum_i \nu_i \mu_i^\#}{T}$$

The total differential of K , since it is a function of both temperature and pressure, is given as:

$$R d \ln K = - \sum_i \nu_i \left[\frac{\partial(\mu_i^\#/T)}{\partial T} dT + \frac{\partial(\mu_i^\#/T)}{\partial P} dP \right]$$

Respecting the functionalities, we can write the above Eq as

$$R d \ln K = - \sum_i \nu_i \left[\frac{\partial(\mu_i^\#/T)}{\partial T} dT + \frac{1}{T} \frac{\partial \mu_i^\#}{\partial P} dP \right] \quad \text{Eq. 6.47}$$


Now, to find the temperature pressure dependence, let us start by a rearranging equation 6.46 as $R \ln K$ equals minus sum over i $\nu_i \mu_i^\#$ by T . The total differential of K , since it is a function of both the temperature, and pressure is given as $R d \ln K$. I am taking the total differential here. R is just a constant. $R d \ln K$ can be written as a function of temperature and a function of pressure. To do that, what I have done here is see here there is a sum ... there is a minus here first, and then there is a sum here. ... Inside the sum, each of these can be ν_i which is not, of course, a function of temperature and pressure. Therefore, that is taken as a constant out here, whereas, $\mu_i^\#$ by T is ... a function of both temperature and pressure.

So, that is written as the total differential here in terms of the partial derivatives $\frac{\partial(\mu_i^\#/T)}{\partial T}$ of $\mu_i^\#$ by T dT plus $\frac{\partial(\mu_i^\#/T)}{\partial P}$ of $\mu_i^\#$ by T dP . Therefore, $R d \ln K$ equals minus of sum over i $\nu_i \frac{\partial(\mu_i^\#/T)}{\partial T} dT$ plus $\frac{1}{T} \frac{\partial \mu_i^\#}{\partial P} dP$. Again respecting the functionalities ... in other words, it means just combining the various terms together, we can write this as $R d \ln K$ equals minus sum over i $\nu_i \frac{\partial(\mu_i^\#/T)}{\partial T} dT$. And, this is not going to vary with pressure. Therefore, you can take $1/T$ out as a constant here. So, $\frac{1}{T} \frac{\partial \mu_i^\#}{\partial P} dP$. Let us call this equation 6.47.

(Refer Slide Time: 06:38)

By following a reasoning similar to the one used to obtain Eq. 6.41, but for solutions, and further, by differentiating μ_i with respect to P instead of T , we can arrive at:

$$R d \ln K = \frac{\Delta H}{T^2} dT - \frac{\Delta V}{T} dP \quad \text{Eq. 6.48}$$

ΔV is the volume change of the reaction mixture due to the reaction



By following a reasoning that is similar to the one used to obtain 6.41, but for solutions ... in other words we had obtained 6.41 for the gas phase reactions. Here, we are going to follow the same reasoning, but for solutions, and further by differentiating μ_i with respect to P instead of T This is what I would like you to do, and the hint here is that you will get it in terms of ΔV , and so on. I will give you about ... since you are already used to it may be about ten minutes. Just look at 6.41 ... go back and look at 6.41. Go a few steps before that; see how we got that. Use the same reasoning to get this particular expression. Ten minutes.

(No audio from 07:29 to 15:33)


You would have obtained $R d \ln K = \frac{\Delta H}{T^2} dT$. Essentially, we have replaced the partial derivatives in terms of the expressions or thermodynamic variables and the relationships we had derived earlier. So, this first derivative gets replaced by $\frac{\Delta H}{T^2} dT$ and the second derivative gets replaced by $-\frac{\Delta V}{T} dP$. Therefore, $R d \ln K = \frac{\Delta H}{T^2} dT - \frac{\Delta V}{T} dP$. Doing such exercises, small exercises, also breaks the monotony of just listening to this and getting actively involved in the process; thereby, the learning is that much better. We will call this equation 6.48 and of course, the ΔV is the volume change of the reaction mixture due to the reaction.

(Refer Slide Time: 16:40)

From Eq. 6.48, we can write:

$$\left(\frac{\partial \ln K}{\partial T}\right)_P = \frac{\Delta H}{RT^2} \quad \text{Eq. 6.49}$$
$$\left(\frac{\partial \ln K}{\partial P}\right)_T = -\frac{\Delta V}{RT} \quad \text{Eq. 6.50}$$

Equations 6.49 and 6.50 can be used to determine the effect of temperature and pressure, respectively, on the equilibrium constant



Usually, the pressure dependence is weak


From equation 6.48 which is this, we can write $\left(\frac{\partial \ln K}{\partial T}\right)_P$ equals $\frac{\Delta H}{RT^2}$ – not quite easy to see that ... we are writing total derivative in terms of the partial derivative. $d \ln K$ equals $\left(\frac{\partial \ln K}{\partial T}\right)_P dT$ plus $\left(\frac{\partial \ln K}{\partial P}\right)_T dP$. So, just comparing the terms here we get $\left(\frac{\partial \ln K}{\partial T}\right)_P$ equals $\frac{\Delta H}{RT^2}$, equation 6.49, and $\left(\frac{\partial \ln K}{\partial P}\right)_T$ equals $-\frac{\Delta V}{RT}$. We will call this equation 6.50.

Equation 6.49 and 6.50 can be used to determine the effect of temperature and pressure on the equilibrium constant. So, that is what we set out to do, and we have done that here. Usually, the pressure dependence is rather weak; rather the pressure dependence of the equilibrium constant is rather weak. It is good to know that also.

(Refer Slide Time: 18:03)

Example 6.1

In a cell undergoing balanced growth at 25 °C, which can be considered being at steady state for short times compared to the times for growth, the concentrations of ATP, ADP, and inorganic phosphate, at a particular time, were found to be 10^{-3} M, 10^{-4} M and 10^{-2} M, respectively. The pH inside the cell can be taken to be 7. The standard $\Delta G'$ value (at physiological conditions) is -7.7 Kcal mol⁻¹. Calculate the ΔG for the hydrolysis of ATP under the above conditions.



Now, we have looked at equilibrium constants of various reactions, the effect of temperature on the gas phase equilibrium constants, the effect of temperature and pressure on the equilibrium constants for reactions that occur in the liquid or solid phases. Now, let us work out an example. There are three examples one after another, deliberately. The reason for having these examples is that these involve clarity in the concepts that you had learnt in the earlier classes also. So, this is some way of refreshing, because you need to know that as a part of thermodynamics of reactions. And therefore, we are going to kind of review that in the context of the problem.

We are going to see and understand better whatever is required to solve these problems. But the point that I am trying to make here is that you will need principles, clarity of principles that you had learnt much earlier in your twelfth standard and previous courses on chemical thermodynamics to be able to do these examples. Because of that I am going to give you significant time to do this before I present the solution. And example 6.1 the first one of the three, we will do one today and may be the other two in the next class. This is in the context of course, bio-systems. In a cell undergoing balanced growth at twenty five degrees C, which can be considered as being at steady state for short times compared to the times of growth. The concentrations of ATP ... recall what ATP is? Adenosine Tri-Phosphate, the energy currency of the cell.

ATP, it is a nucleotide, if you recall the type of molecule – ADP, adenosine di-phosphate and inorganic phosphate. I am sure, now you recall the relationship between ATP, ADP and inorganic phosphate from your biochemistry course. The concentrations of ATP,

ADP and inorganic phosphate, P_i , at a particular time were found to be ten power minus three molar, ten power minus four molar and ten power minus two molar, respectively. The pH inside the cell can be taken to be 7, this is a reasonably good assumption.

For example, the pH inside a mammalian cell varies anywhere from about 7.2 to 7.3. Typically, the pH is very well regulated inside the cell, except if there is a drastic change in the function of the cell. For example, when the cell goes into hibernation, which is a very drastic change, then there is a change of about 0.8 to 1 unit in pH. For metabolic activities, there is a change of about 0.2 units, and so on and so forth, as the activity proceeds. And sometimes, it is as high as about 0.5 units. But, for the purposes of this particular problem we will consider the pH inside the cell as a constant.

The standard ΔG° value at physiological conditions, there is a whole meaning attached to this ΔG° value. It is not the same as ΔG° value. I hope you recall that at physiological conditions that it is ΔG° . What is physiological about it? You must be able to recall; it is minus 7.7 kilocalories per mole, and calculate the ΔG , the Gibbs free energy change for the hydrolysis of ATP under the above conditions. What I would like you to do is take about twenty minutes to this. It is going to take your time, first to recall whatever is required. It is quite a simple calculation, if you know what is needed. But it is going to take your time to recall those – go back to your notes, check that of the previous courses and so on. So, take about twenty minutes or maybe even twenty, twenty five minutes and then come back with the solution. I will present the detailed solution after you come back. Go ahead, please.

(No audio from 22:35 to 42:41)

You would have recalled quite a few principles from the earlier course. Let us see what these are or what is required to solve this particular example.


(Refer Slide Time: 42:55)

Solution

From earlier courses, you would know that for a reaction

$$\Delta G = \Delta G^{\circ} + R T \ln \left(\frac{\prod (\text{product conc})^{\text{stoichiometric coefficient}}}{\prod (\text{reactant conc})^{\text{stoichiometric coefficient}}} \right)$$
$$= \Delta G^{\circ} + R T \ln \left(\prod_i c_i^{\nu_i} \right)$$

Recall that ΔG° is defined at standard conditions of T (25 °C) and P (1 atm) and when all the concentrations (products and reactants) are each at 1 M



From earlier courses, we know that for a reaction, ΔG equals ΔG° plus $RT \ln$ of the product of the product concentrations raised to the power of the appropriate stoichiometric coefficient or the relevant stoichiometric coefficient, divided by the product of the reactant concentrations, each raised to its own stoichiometric coefficient. In other words, ΔG equals ΔG° plus $RT \ln$ product of $C_i^{\nu_i}$. As you can recognize now, our convention of ν_i being positive for products and negative for reactants automatically takes care of this particular formulation here.

Now, recall that ΔG° is defined at standard conditions of temperature, which is twenty five degree C and pressure, which is one atmosphere. And, more importantly in the context of whatever we are going to discuss now, when all the concentrations that is the concentrations of products as well as reactants are each at one molar concentration. That is how ΔG° is defined. But what is the problem with this? It is perfectly fine for chemical reactions, but what is the problem with this for biological reactions? Would you recall?

(Refer Slide Time: 44:32)

When the H^+ ion is involved in the reaction, e.g. in the reaction under consideration, it becomes difficult, because for the standard conditions, its concentration should also be set to 1 M, which corresponds to a pH of 0

The pH of 0 is non-physiological since the proteins and enzymes in the cell would be deactivated at that pH

Thus, the standard conditions of relevance to Biological systems are when the pH equals 7, or the H^+ ion concentration equals $10^{-7} M$

The temperature, pressure and the 1 M concentration requirement of the other reactants and products remain the same

NPTEL

The problem comes about whenever hydrogen ion is involved in a reaction, which is pretty much most of the time, because most reactions that occur in a cell are ionic reactions. So, when the hydrogen ion is involved in the reaction it becomes difficult, because for the standard conditions its concentration should also be set to one molar. And, what happens when you set the concentration of hydrogen ions to one molar, the pH which is nothing but ... the negative log of the hydrogen ion concentration as a first approximation, turns out to be zero.

... The cell is completely gone at a pH of zero, because of the cell typically works at a pH of seven or around seven. And, pH of zero is non-physiological since the proteins and the enzymes in the cell would certainly be deactivated at that pH. And therefore, people came up with a different standard or a different set of standard conditions of relevance to biological systems. ... That is, the pH is taken to be equal to seven or the hydrogen ion concentration is taken to be equal to ten power minus seven molar.


Whereas, the other conditions the other standard conditions, the temperature, twenty five degree C, pressure, one atmosphere, and the one molar concentration requirement for all the other reactants and products remain the same. I hope you recall this now. Therefore, you know, this and the hydrogen ion concentration being ten power minus seven molar is the set of standard conditions for biological reactions, which is different from the standard conditions for normal reactions.

(Refer Slide Time: 46:26)

Under such conditions, $\Delta G'$ is defined, in which H^+ concentration is not taken into account in the definition, explicitly, as long as it is understood that the pH is 7. Thus,

$$\Delta G = \Delta G' + RT \ln \left(\prod_i c_i^{\nu_i} \right)$$

For the current reaction

$$ATP + H_2O \leftrightarrow ADP + P_i \quad \rightleftharpoons$$


And therefore, we use a different terminology also. Before that, I should say that under such conditions, delta G dash – this is the different terminology that I am talking about – delta G dash is defined instead of delta G naught, we use a delta G dash. This is defined in which the hydrogen ion concentration is not taken into account in the definition explicitly, as long as it is understood that the pH is seven. It is essentially saying that the hydrogen ion concentration is taken as ten power minus seven molar.


And therefore, for our biological reactions, the equation of relevance to use is delta G equals delta G dash plus RT ln ... the product over all i or C i power mu i. And for the current reaction, which is ATP plus water giving you ADP plus P i – this is the reaction that we are considering here ATP, ADP and P i, ... the concentrations of which we knew earlier.

(Refer Slide Time: 47:28)

Under excess water conditions, when the concentration of water is taken to be a constant, we can write

$$\Delta G = \Delta G' + R T \ln \left(\frac{[ADP] [P_i]}{[ATP]} \right)$$

Thus,

$$\Delta G = -7700 + 8.31 \times 298 \ln \left(\frac{[10^{-4}] [10^{-2}]}{[10^{-3}]} \right)$$
$$= -11.79 \text{ Kcal mol}^{-1}$$


Under excess water conditions, when the concentration of water can be taken to be a constant, I hope you recall this trick that we use: we can say that delta G equals delta G dash plus RT ln concentration of the products ... this is hydrolysis of ATP ... so, concentration of the product, ADP, and concentration of the other product, P i, divided by the concentration of ATP. Under the snapshot conditions, that were given delta G equals delta G dash which is minus 7.7 kilo calories per mole or minus seventy seven hundred calories per mole plus 8.31 times 298, temperature, and all these concentrations taken in molar quantities. That turns out to be ... the R has to be in calories, it is not 8.31 it is 1.983. This will turn out to be minus 11.79 kilo calories per mole.

I think we are out of time. I think we need to redo this particular calculation, why do not you redo this particular calculation and actually tell me, because I have taken this to be in kilo calories I should also take the relevant set of units here; just check and tell me whether this value is correct. We will stop here for now. When we come back in the next class, we will work out the other two problems that I mentioned, which also will involve whatever we did in this particular module plus a lot of background information that you need to brush up.